

Updated Status of Federally Listed ESUs of West Coast Salmon and Steelhead

West Coast Salmon Biological Review Team

Northwest Fisheries Science Center

2725 Montlake Boulevard East
Seattle, WA 98112

Southwest Fisheries Science Center

Santa Cruz Laboratory
110 Shaffer Road
Santa Cruz, CA 95060

July 2003

This is a scientific report. It does not represent or set policy for NMFS or NOAA, regardless of whether any statements contained herein may appear to do so.

A.2.8 SACRAMENTO RIVER WINTER-RUN CHINOOK SALMON

**Primary contributor: Steven T. Lindley
(Southwest Fisheries Science Center – Santa Cruz Lab)**

A.2.8.1 Summary of Previous BRT Conclusions

The status of chinook salmon coastwide was formally assessed in 1998 (Myers et al. 1998); however, NMFS had previously recognized Sacramento River winter-run chinook as a “distinct population segment” under the ESA (NMFS 1987).

Summary of major risk factors and status indicators

Historically, winter-run chinook salmon were dependent on access to spring-fed tributaries to the upper Sacramento River that stayed cool during the summer and early fall. Adults enter freshwater in early winter and spawn in the spring and summer. Juveniles rear near the spawning location until at least the fall, when water temperatures in lower reaches are suitable for migration. Winter-run chinook salmon were abundant and comprised populations in the McCloud, Pit, and Little Sacramento, with perhaps smaller populations in Battle Creek and the Calaveras River. On the basis of commercial fishery landings in the 1870s, Fisher (1994) estimated that the total run size of winter-run chinook salmon may have been 200,000 fish.

The most obvious challenge to winter-run chinook salmon was the construction of Shasta Dam, which blocked access to the entire historic spawning habitat. It was not expected that winter-run chinook salmon would survive this habitat alteration (Moffett 1949). Cold-water releases from Shasta, however, created conditions suitable for winter-run chinook salmon for roughly 100 km downstream from the dam. Presumably, there were several independent populations of winter-run chinook salmon in the Pitt, McCloud, and Little Sacramento Rivers, and various tributaries to these rivers, such as Hat Creek and the Fall River. These populations merged to form the present single population. If there ever were populations in Battle Creek and the Calaveras River, they have been extirpated.

In addition to having only a single extant population dependent on artificially created conditions, winter-run chinook salmon face numerous other threats. Chief among these is small population size—escapement fell below 200 fish in the 1990s. Population size declined monotonically from highs of near 100,000 fish in the late 1960s, indicating a sustained period of poor survival. There are questions of genetic integrity due to winter-run chinook salmon having passed through several bottlenecks in the 20th century. Other threats include inadequately screened water diversions, predation at artificial structures and by non-native species, pollution from Iron Mountain Mine (among other sources), adverse flow conditions, high summer water temperatures, unsustainable harvest rates, passage problems at various structures (e.g., Red Bluff Diversion Dam), and vulnerability to drought.

Previous BRT conclusions

The chinook salmon BRT spent little time considering the status of winter-run chinook salmon, because winter-run chinook salmon were already listed as endangered at the time of previous BRT meetings.

Listing status

Winter-run chinook salmon were listed as Threatened in 1990 and reclassified as Endangered 1994.

A.2.8.2 New Data and Updated Analyses

Viability assessments

Two studies have been done on the population viability of Sacramento River winter-run chinook salmon. Botsford and Brittnacher (1998), in a paper that is part of the draft recovery plan, developed de-listing criteria using a simple age-structured, density-independent model of spawning escapement. They concluded, on the basis of the 1967-1995 data, that winter-run chinook salmon were certain to fall below the quasi-extinction threshold of three consecutive spawning runs with less than 50 females.

Lindley and Mohr (2003) developed a slightly more complex Bayesian model of winter-run chinook salmon spawning escapement that allowed for density dependence and a change in population growth rate in response to conservation measures initiated in 1989. This model, due to its allowance for the growth rate change, its accounting for parameter uncertainty, and use of newer data (through 1998), suggested a lower but still biologically significant expected quasi-extinction probability of 28%.

Draft recovery plan

The draft recovery plan for winter-run chinook salmon (NMFS 1997) provides a comprehensive review of the status, life history, habitat requirements, and risk factors of winter-run chinook salmon. It also provides a recovery goal: an average of 10,000 females spawners per year and a $\lambda \geq 1.0$ calculated over 13 years of data (assuming a certain level of precision in spawning escapement estimates).

New abundance data

The winter-run chinook salmon spawning run has been counted at Red Bluff Diversion Dam (RBDD) fish ladders since 1967. Escapement has been estimated with a carcass survey since 1996. Through the mid-1980s, the RBDD counts were very reliable. At that time, changes to the dam operation were made to alleviate juvenile and adult passage problems. Now, only the tail end of the run (about 15% on average) is forced over the ladders, greatly reducing the accuracy of the RBDD counts. The carcass mark-recapture surveys were initiated to improve

escapement estimates. The two measures are in very rough agreement, and there are substantial problems with both estimates, making it difficult to choose one as more reliable than the other. One problem with the carcass-based estimate is the estimation of the probability of capturing carcasses—it appears that the probability of initial carcass recovery depends strongly on the sex of the fish, the size of the fish, and possibly on whether it has been previously recovered. In the winter-run chinook salmon carcass surveys, a high ratio of female to males is observed (e.g., Snider et al. 1999), and several studies of salmon carcass recovery have noted that females are recovered with a higher probability than males, presumably because of the different behavior of males and females (e.g., Shardlow et al. 1986 and references therein). In spite of these problems, both abundance measures suggest that the abundance of winter-run chinook salmon is increasing. Based on the RBDD counts, the winter-run chinook salmon population has been growing rapidly since the early 1990s (Figure A.2.8.1), with a short-term trend of 0.26 (Table A.2.8.1). On the population growth rate-population size space, the winter-run chinook salmon population has a somewhat low population growth and moderate size compared to other Central Valley salmonid populations (Figure A.2.8.2).

Table A.2.8.1. Summary statistics for trend analyses. Numbers in parentheses are 0.90 confidence intervals. Results for other populations are shown for comparison.

Population	5-yr mean	5-yr min	5-yr max	λ	μ	LT trend	ST trend
Sacramento River winter-run chinook	2,191	364	65,683	0.97 (0.87, 1.09)	-0.10 (-0.21, 0.01)	-0.14 (-0.19, -0.09)	0.26 (0.04, 0.48)
Butte Creek spring-run chinook	4,513	67	4,513	1.30 (1.09, 1.60)	0.11 (-0.05, 0.28)	0.11 (0.03, 0.19)	0.36 (0.03, 0.70)
Deer Creek spring-run chinook	1,076	243	1,076	1.17 (1.04, 1.35)	0.12 (-0.02, 0.25)	0.11 (0.02, 0.21)	0.16 (-0.01, 0.33)
Mill Creek spring-run chinook	491	203	491	1.19 (1.00, 1.47)	0.09 (-0.07, 0.26)	0.06 (-0.04, 0.16)	0.13 (-0.07, 0.34)
Sacramento River steelhead	1,952	1,425	12,320	0.95 (0.90, 1.02)	-0.07 (-0.13, 0.00)	-0.09 (-0.13, -0.06)	NA

Winter-run chinook salmon may be responding to a number of factors, including wetter-than-normal winters, changes in ocean harvest regulations since 1995 significantly reducing harvest, changes in RBDD operation, improved temperature management on the Upper Sacramento (including installation of a cold-water release device on Shasta Dam), water quality improvements due to remediation of Iron Mountain Mine discharges, changes in operations of the state and federal water projects, and a variety of other habitat improvements. While the status of winter-run chinook salmon is improving, there is only one winter-run chinook salmon population and it is dependent on cold-water releases of Shasta Dam, which could be vulnerable to a prolonged drought. The recent 5-year geometric mean is only 3% of the maximum post-1967 5-year geometric mean.

The RBDD counts are suitable for modeling as a random-walk-with-drift (also known as the “Dennis model” [Dennis et al. 1991]). In the RWWD model, population growth is described by exponential growth or decline:

$$N_{t+1} = N_t \exp(\mu + \eta_t), \quad (1)$$

where N_t is the population size at time t , μ is the mean population growth rate, and η_t is a normal random variable with mean=0 and variance = σ_p^2 .

Table A.2.8.2. Parameter estimates for the constant-growth and step-change models applied to winter-run chinook salmon. Numbers in parentheses indicate 90% confidence intervals.

parameter	Model	
	constant μ	step change μ
μ	-0.085 (-0.181, 0.016)	-0.214 (-0.322, -0.113)
δ	NA	0.389 (0.210, 0.574)
σ_p^2	0.105 (0.094, 0.122)	0.056 (0.046, 0.091)
σ_m^2	0.0025 (2.45E-6, 0.0126)	0.011 (3.92E-6, 0.022)
$P_{100}(\text{ext})^{[a]}$	0.40 (0.00, 0.99)	0.003 (0.0, 0.0)

^[a] Probability of extinction (pop. size < 1 fish) within 100 years.

The RWWD model, as written in Equation 1, ignores measurement error. Observations (y_t) can be modeled separately,

$$y_t = N_t \exp(\varepsilon_t), \quad (2)$$

where ε_t is a normal random variable with mean = 0 and variance = σ_m^2 . Equations 1 and 2 together define a state-space model that, after linearizing by taking logarithms, can be estimated using the Kalman filter (Lindley in press).

A recent analysis of the RBDD data (Lindley and Mohr 2003) indicated that the population growth since 1989 was higher than in the preceding period. For this reason, I fit two forms of the RWWD model—one with a fixed growth rate (constant-growth model) and another with a growth rate with a step-change in 1989, when conservation actions began (step-change model, $\mu_t = \mu$ for $t < 1989$, $\mu_t = \mu + \delta$ for $t \geq 1989$). In both cases, a 4-year running sum was applied to the spawning escapement data to form a total population estimate (Holmes 2001). Results of model fitting are shown in Table A.2.8.2. The constant-growth model satisfies all model diagnostics, although visual inspection of the residuals shows a strong tendency to under-predict abundance in the most recent 10 years. The residuals of the step-change model fail the Shapiro-Wilks test for normality; the residuals look truncated on the positive side, meaning that good years are not as extreme as bad years. Winter-run chinook salmon growth rate might be better modeled as a

mixture between a normal distribution and another distribution reflecting near-catastrophic population declines caused by episodic droughts.

According to Akaike's information criterion (AIC), the step-change model is a much better approximation to the data than the constant population growth rate model, with an AIC difference of 9.61 between the two models (indicating that the data provide almost no support for the constant-growth model). The step-change model suggests the winter-run chinook salmon population currently has a λ of 1.21, while for the constant population growth rate model, $\lambda = 0.97^5$. The extinction risks predicted by the two models are extremely different: winter-run chinook salmon have almost no risk of extinction if the apparent recent increase in λ holds in the future, but are certain to go extinct if the population grows at its average rate, with a most likely time of extinction being 100 years. While it would be dangerous to assume that recent population growth will hold indefinitely, it does appear that the status of winter-run chinook salmon is improving.

Harvest impacts

Substantial changes in ocean fisheries off central and northern California have occurred since the last status review (PFMC 2002a, b). Ocean harvest rate of winter-run chinook salmon is thought to be a function of the Central Valley chinook salmon ocean harvest index (CVI), which is defined as the ratio of ocean catch south of Point Arena to the sum of this catch and the escapement of chinook salmon to Central Valley streams and hatcheries. Note that other stocks (e.g., Klamath chinook salmon) contribute to the catch south of Point Arena, and that fish from the Central Valley are caught in Oregon fisheries. This harvest index ranged from 0.55 to nearly 0.80 from 1970 to 1995, when harvest regimes were adjusted to protect winter-run chinook salmon. In 2001, the CVI fell to 0.27. The reduction in harvest is presumably at least partly responsible for the record spawning escapement of fall-run chinook salmon ($\approx 540,000$ fish in 2001) and concurrent increases in other chinook salmon runs in the Central Valley.

Because they mature before the ocean fishing season, winter-run chinook salmon should have lower harvest rates than fall-run chinook salmon, if they have similar age-at-maturity. At the time of the last status review, the only information on the harvest rate of winter-run chinook salmon came from a study conducted in the 1970s. Hallock and Fisher (1985) reported that the average catch/(catch+escapement) for the 1969-71 broodyears was 0.40 for the ocean fishery. For the 1968-1975 period, freshwater sport fisheries caught an average of 10% of the winter chinook salmon run.

The recent release of significant numbers of ad-clipped winter-run chinook salmon provides new, but limited, information on the harvest of winter-run chinook salmon in coastal recreational and troll fisheries. The Pacific Fisheries Management Council's Sacramento River Winter and Spring Chinook salmon Workgroup (SRWSCW) conducted a cohort reconstruction of the 1998 broodyear (PFMC 2003). Winter-run chinook salmon are mainly vulnerable to ocean fisheries as 3-year olds. SRWSCW calculated, on the basis of 123 coded-wire-tag

⁵In this section of the document, λ is defined as $\exp(\mu + \sigma_p^2 / 2)$, the *mean* annual population growth rate.

recoveries, that the ocean fishery impact rate on 3-year-olds was 0.23, and the in-river sport fishery impact rate was 0.24. These impacts combine to reduce escapement to $100 \times (1 - 0.23) \times (1 - 0.24) = 59\%$ of what it would have been in the absence of fisheries, assuming no natural mortality during the fishing season. The high estimated rate of harvest in the river sport fishery, which arises from the recovery of eight coded-wire tags, was a surprise because salmon fishing is closed from January 15 to July 31 to protect winter-run chinook salmon. The tags were recovered in late December/early January, at the tail end of the fishery for late-fall-run chinook salmon. The estimate of river sport fishery impact is much less certain than the ocean fishery impact estimate because of the lower number of tag recoveries, less rigorous tag sampling, and larger expansion factors. The California Fish and Game Commission is moving forward with an emergency action to amend sport fishing regulations to ban retention of salmon caught in river sport fisheries on January 1 rather than January 15. Had such regulations been in place in 1999/2000, the freshwater harvest rate would have been 20% of that observed.

New hatchery information

Livingston Stone National Fish Hatchery (LSNFH) was constructed at the base of Shasta Dam in 1997, with the sole purpose of helping to restore natural production of winter-run chinook salmon. LSNFH was designed as a conservation hatchery with features intended to overcome the problems of CNFH (better summer water quality, natal water source). All production is ad-clipped. Each individual considered for use as broodstock is genotyped to ensure that it is a winter-run chinook salmon. No more than 10% of the broodstock is composed of hatchery-origin fish, and no more than 15% of the run is taken for broodstock, with a maximum of 120 fish. Figure 3 shows the number of winter-run chinook salmon released by CNFH/LSNFH; Figure 4 shows the number of winter-run chinook salmon spawners taken into the hatchery.

A.2.8.3 New Comments

The California State Water Contractors, the San Luis and Delta-Mendota Water Authority, and the Westlands Water District recommend that the listing status of winter-run chinook salmon be changed from Endangered to Threatened. They base this proposal on the recent upturn of adult abundance, recently initiated conservation actions (restoration of Battle Creek, ocean harvest reductions, screening of water diversions, remediation of Iron Mountain Mine, and improved temperature control), and a putative shift in ocean climate in 1999.




Figure A.2.8.1. Estimated winter-run chinook spawner abundance as determined by RBDD fish ladder (solid line) and carcass mark-recapture (dashed line).




Figure A.2.8.2. Abundance and growth rate of Central Valley salmonid populations. Open circle- steelhead; open squares- spring chinook; filled triangle- winter-run chinook; small black dots- other chinook stocks. Error bars represent central 0.90 probability intervals for μ estimates. (Note: as defined in other sections of the status reviews, $\mu \approx \log(\lambda)$.)




Figure A.2.8.3. Number of juvenile winter-run chinook released by Coleman and Livingston Stone National Fish Hatcheries.




Figure A.2.8.4. Number of adult winter-run chinook collected for broodstock by Coleman and Livingston Stone National Fish Hatcheries.

A.3 CHINOOK SALMON BRT CONCLUSIONS

Snake River fall-run chinook salmon ESU

A majority (60%) of the BRT votes for this ESU fell in the “likely to become endangered” category, with minorities falling in the “danger of extinction” and “not likely to become endangered” categories (Table A.3.1). This represented a somewhat more optimistic assessment of the status of this ESU than was the case at the time of the original status review, when the BRT concluded that Snake River fall-run chinook salmon “face a substantial risk of extinction if present conditions continue” (Waples et al. 1991). The BRT found moderately high risks in all VSP elements, with mean risk matrix scores ranging from 3.0 for growth rate/productivity to 3.6 for spatial structure (Table A.3.2).

On the positive side, the number of natural origin spawners in 2001 was well in excess of 1000 for the first time since counts at Lower Granite Dam began in 1975. Management actions have reduced (but not eliminated) the fraction of fish passing Lower Granite Dam that are strays from out-of-ESU hatchery programs. Returns in the last two years also reflect an increasing contribution from supplementation programs based on the native Lyons Ferry Broodstock. With the exception of the increase in 2001, the ESU has fluctuated between approximately 500-1000 adults, suggesting a somewhat higher degree of stability in growth rate and trends than is seen in many other salmon populations.

In spite of the recent increases, however, the recent geometric mean number of naturally produced spawners is still less than 1000, a very low number for an entire ESU. Because of the large fraction of naturally spawning hatchery fish, it is difficult to assess the productivity of the natural population. The relatively high risk matrix scores for spatial structure and diversity (3.5-3.6) reflect the concerns of the BRT that a large fraction of historic habitat for this ESU is inaccessible, diversity associated with those populations has been lost, the single remaining population is vulnerable to variable environmental conditions or catastrophes, and continuing immigration from outside the ESU at levels that are higher than occurred historically. Some BRT members were concerned that the efforts to remove stray, out-of-ESU hatchery fish only occur at Lower Granite Dam, well upstream of the geographic boundary of this ESU. Specific concerns are that natural spawners in lower river areas will be heavily affected by strays from Columbia River hatchery programs, and that this approach effectively removes the natural buffer zone between the Snake River ESU and Columbia River ocean-type chinook salmon. The effects of these factors on ESU viability are not known, as the extent of natural spawning in areas below Lower Granite Dam is not well understood, except in the lower Tucannon River.

Snake River spring/summer-run chinook salmon ESU

About two-thirds (68%) of the BRT votes for this ESU fell in the “likely to become endangered” category, with minorities falling in the “danger of extinction” and “not likely to become endangered” categories (Table A.3.1). As indicated by mean risk matrix scores, the BRT had much higher concerns about abundance (3.6) and growth rate/productivity (3.5) than for spatial structure (2.2) and diversity (2.3) (Table A.3.2).

Although there are concerns about loss of an unquantified number of spawning aggregations that historically may have provided connectivity between headwater populations, natural spawning in this ESU still occurs in a wide range of locations and habitat types.

Like many others, this ESU saw a large increase in escapement in many (but not all) populations in 2001. The BRT considered this an encouraging sign, particularly given the record low returns seen in many of these populations in the mid 1990s. However, recent abundance in this ESU is still short of the levels that the proposed recovery plan for Snake River salmon indicated should be met over at least an eight year period (NMFS 1995). The BRT considered it a positive sign that the non-native Rapid River broodstock has been phased out of the Grande Ronde system, but the relatively high level of both production/mitigation and supplementation hatcheries in this ESU leads to ongoing risks to natural populations and makes it difficult to assess trends in natural productivity and growth rate.

Upper Columbia River spring-run chinook salmon ESU

Assessments by the BRT of the overall risks faced by this ESU were divided, with a slight majority (53%) of the votes being cast in the “danger of extinction” category and a substantial minority (45%) in the “likely to be endangered” category (Table A.3.1). The mean risk matrix scores reflect strong ongoing concerns regarding abundance (4.4) and growth rate/productivity (4.5) in this ESU and somewhat less (but still significant) concerns for spatial structure (2.9) and diversity (3.5) (Table A.3.2).

Many populations in this ESU have rebounded somewhat from the critically low levels that immediately preceded the last status review evaluation, and this was reflected in the substantial minority of BRT votes cast that were not cast in the “danger of extinction” category. Although this was considered an encouraging sign by the BRT, the last year or two of higher returns come on the heels of a decade or more of steep declines to all time record low escapements. In addition, this ESU continues to have a very large influence by hatchery production, both from production/mitigation and supplementation programs. The extreme management measures taken in an effort to maintain populations in this ESU during some years in the late 1990s (collecting all adults from major basins at downstream dams) are a strong indication of the ongoing risks to this ESU, although the associated hatchery programs may ultimately play a role in helping to restore self-sustaining natural populations.

Lower Columbia River chinook salmon ESU

A majority (71%) of the BRT votes for this ESU fell in the “likely to become endangered” category, with minorities falling in the “danger of extinction” and “not likely to become endangered” categories (Table A.3.1). Moderately high concerns for all VSP elements are indicated by mean risk matrix scores ranging from 3.2 for abundance to 3.9 for diversity (Table A.3.2).

All of the risk factors identified in previous reviews were still considered important by the BRT. The Willamette/Lower Columbia River TRT has estimated that 8-10 historic populations

in this ESU have been extirpated, most of them spring-run populations. Near loss of that important life history type remains in important BRT concern. Although some natural production currently occurs in 20 or so populations, only one exceeds 1000 spawners. High hatchery production continues to pose genetic and ecological risks to natural populations and to mask their performance. Most populations in this ESU have not seen as pronounced increases in recent years as occurred in many other geographic areas.

Upper Willamette River chinook salmon ESU

A majority (70%) of the BRT votes for this ESU fell in the “likely to become endangered” category, with minorities falling in the “danger of extinction” and “not likely to become endangered” categories (Table A.3.1). The BRT found moderately high risks in all VSP elements (mean risk matrix scores ranged from 3.1 for growth rate/productivity to 3.6 for spatial structure) (Table A.3.2).

Although the number of adult spring-run chinook salmon crossing Willamette Falls is in the same range (about 20,000–70,000) it has been for the last 50 years, a large fraction of these are hatchery produced. The score for spatial structure reflects concern by the BRT that perhaps a third of the historic habitat used by fish in this ESU is currently inaccessible behind dams, and the BRT remained concerned that natural production in this ESU is restricted to a very few areas. Increases in the last 3-4 years in natural production in the largest remaining population (the McKenzie) were considered encouraging by the BRT. With the relatively large incidence of hatchery fish, it is difficult to determine trends in natural production.

Puget Sound chinook salmon ESU

A majority (74%) of the BRT votes for this ESU fell in the “likely to become endangered” category, with minorities falling in the “danger of extinction” and “not likely to become endangered” categories (Table A.3.1). The BRT found moderately high risks in all VSP elements, with mean risk matrix scores ranging from 2.9 for spatial structure to 3.6 for growth rate/productivity (Table A.3.2).

Most population indices for this ESU have not changed substantially since the last BRT assessment. The Puget Sound TRT has identified approximately 31 historic populations, of which 9 are believed to be extinct, with most of the populations that have been lost being early run. Other concerns noted by the BRT are the concentration of the majority of natural production in just two basins, high levels of hatchery production in many areas of the ESU, and widespread loss of estuary and lower floodplain habitat diversity (and, likely, associated life history types). Although populations in this ESU have not experienced the sharp increases in the last 2-3 years seen in many other ESUs, more populations increased than decreased over the 4 years since the last BRT assessment. After adjusting for changes in harvest rates, however, trends in productivity are less favorable. Most populations are relatively small, and recent natural production within the ESU is only a fraction of estimated historic run size. On the positive side, harvest rates for all populations have been reduced from their peaks in the 1980s, and some hatchery reforms have been implemented (e.g., elimination of many net pen programs that were leading to widespread straying, and transition of other programs to more local

broodstocks). The BRT felt that these management changes should help facilitate recovery if other limiting factors (especially habitat degradation) are also addressed. The BRT felt that the large recovery effort organized around the Puget Sound Shared Strategy was a positive step because it could help to link and coordinate efforts in many separate, local watersheds.

California Coastal chinook salmon ESU

A majority (67%) of the BRT votes for this ESU fell in the “likely to become endangered” category, with votes falling in the “danger of extinction” category outnumbering those in “not warranted” category by nearly 2-to-1 (Table A.3.1). The BRT found moderately high risks in all VSP elements, with mean risk matrix scores ranging from 3.1 for diversity to 3.9 for abundance (Table A.3.2).

The BRT was concerned by continued evidence of low population sizes relative to historical abundance and mixed trends in the few time series of abundance indices available for analysis, and by the low abundances and potential extirpations of populations in the southern part of the ESU. The BRT’s concerns regarding genetic integrity of this ESU were moderate or low relative to similar issues for other ESUs because 1) hatchery production in this ESU is on a minor scale, and 2) current hatchery programs are largely focused on supplementing and restoring local populations. However, the BRT did have concerns with respect to diversity that were based largely on the loss of spring-run chinook salmon in the Eel River basin and elsewhere in the ESU, and to a lesser degree on the potential loss of diversity concurrent with low abundance or extirpation of populations in the southern portion of the ESU. Overall, the BRT was strongly concerned by the paucity of information and resultant uncertainty associated with estimates of abundance, natural productivity and distribution of chinook salmon in this ESU.

Sacramento River winter-run chinook salmon ESU

A majority (60%) of the BRT votes fell into the “in danger of extinction” category, with a minority (38%) voting for the “likely to become endangered” and only 2% voting for “not warranted.” (Table A.3.1). The main VSP concerns were in the spatial structure and diversity categories (4.8 and 4.2, respectively), although there was significant concern in the abundance and productivity categories (3.7 and 3.5, respectively) (Table A.3.2).

The main concerns of the BRT relate to the lack of diversity within this ESU. The BRT was very troubled by the fact that this ESU is represented by a single population that has been displaced from its historic spawning habitat into an artificial habitat created and maintained by a dam. The BRT presumed that several independent populations of winter-run chinook salmon were merged into a single population, with the potential for a significant loss of life history and genetic diversity. Furthermore, the population has passed through at least two recent bottlenecks—one when Shasta Dam was filled and another in the late 1980s-early 1990s—that probably further reduced genetic diversity. The population has been removed from the environment where it evolved, dimming its long-term prospects for survival. The BRT was modestly heartened by the increase in abundance since the lows of the late 1980s and early 1990s.

Central Valley spring-run chinook salmon ESU

A large majority (69%) of the BRT votes fell into the “likely to become endangered” category, with a minority (27%) of votes going to “in danger of extinction” and 4% “not warranted” (Table A.3.1). There was roughly equal concern about abundance, spatial structure and diversity (3.5-3.8), and less concern about productivity (2.8) (Table A.3.2).

A major concern of the BRT was the loss of diversity caused by the extirpation of spring-run chinook salmon populations from most of the Central Valley, including all San Joaquin tributaries. The only populations left in the Sierra Nevada ecoregion are supported by the Feather River hatchery. Another major concern of the BRT was the small number and location of extant spring-run chinook salmon populations-- only three streams, originating in the southern Cascades, support self-sustaining runs of spring-run chinook salmon, and these three streams are close together, increasing their vulnerability to catastrophe. Two of the three extant populations are fairly small, and all were recently quite small. The BRT was also concerned about the Feather River spring-run chinook salmon hatchery population, which is not in the ESU but does produce fish that potentially could interact with other spring-run chinook salmon populations, especially given the off-site release of the production.

Table A.3.1. Tally of FEMAT vote distribution regarding the status of 9 chinook salmon ESUs reviewed by the chinook salmon BRT. Each of 15 BRT members allocated 10 points among the three status categories.

ESU	At Risk of Extinction	Likely to Become Endangered	Not Likely to Become Endangered
Snake River fall-run	38	91	21
Snake River spring/summer-run	30	102	18
Upper Columbia River spring-run	79	67	4
Puget Sound	12	111	27
Lower Columbia River	25	107	18
Upper Willamette River	32	105	13
California Coastal ¹	36	100	13
Sacramento River winter-run ²	78	49	3
CA Central Valley spring-run ²	35	90	5

¹One BRT member assigned 9 points

²Votes tallied for 13 BRT members

Table A.3.2. Summary of risk scores (1 = low to 5 = high) for four VSP categories (see section "Factors Considered in Status Assessments" for a description of the risk categories) for the 9 chinook salmon ESUs reviewed. Data presented are means (range).

ESU	Abundance	Growth Rate/Productivity	Spatial Structure and Connectivity	Diversity
Snake River fall-run	3.4 (2-5)	3.0 (2-5)	3.6 (2-5)	3.5 (2-5)
Snake River spring/summer-run	3.6 (2-5)	3.5 (3-5)	2.2 (1-3)	2.3 (1-3)
Upper Columbia River spring-run	4.4 (3-5)	4.5 (3-5)	2.9 (2-4)	3.5 (2-5)
Puget Sound	3.3 (2-4)	3.6 (3-4)	2.9 (2-4)	3.2 (2-4)
Lower Columbia River	3.2 (2-4)	3.7 (3-5)	3.5 (3-4)	3.9 (3-5)
Upper Willamette River	3.7 (2-5)	3.1 (2-5)	3.6 (3-4)	3.2 (2-4)
California Coastal ¹	3.9 (3-5)	3.3 (3-4)	3.2 (2-4)	3.1 (2-4)
Sacramento River winter-run ²	3.7 (3-5)	3.5 (2-5)	4.8 (4-5)	4.2 (3-5)
CA Central Valley spring-run ²	3.5 (3-4)	2.8 (2-4)	3.8 (3-5)	3.8 (3-5)

A.4 REFERENCES

- Anonymous. 1998. ODFW Columbia River Management unpublished files: Willamette Falls fish passage 1946-97.
- Beamesderfer, R. C. P., H. P. Schaller, M. P. Zimmerman, C. E. Petrosky, O. P. Langness, and L. LaVoy. 1997. Spawner-recruit data for spring and summer chinook salmon populations in Idaho, Oregon, and Washington. Unpublished draft manuscript dated 6/23/97, 78 p. (Compiled and edited by ESSA Technologies Ltd., Vancouver, B.C.)
- Bennett, D. E. 1986. Fish passage at Willamette Falls in 1985, Annual report, January 1-December 31, 1985. ODFW, Portland, OR.
- Bennett, D. E. and C.A. Foster. 1990. 1989 Willamette River spring chinook run, fisheries, and passage at Willamette Falls. ODFW.
- Bennett, D. E. and C. A. Foster. 1994. 1993 Willamette River spring chinook salmon run. ODFW, Portland, OR.
- Bennett, D. E. and C.A. Foster. 1995. 1994 Willamette river spring chinook run, fisheries, and passage at Willamette Falls, draft. Unpublished.
- Bishop, S. and A. Morgan (eds.). 1996. Critical habitat issues by basin for natural chinook salmon stocks in the coastal and Puget Sound areas of Washington State. Northwest Indian Fisheries Commission, Olympia, WA, 105 p.
- Botsford, L. W. and J. G. Brittnacher. 1998. Viability of Sacramento River winter-run chinook salmon. *Conserv. Biol.* 12:65-79.
- Bugert, R. M., and B. Hopley. 1989. The Snake River fall chinook salmon egg bank program: The final chapter. Washington Department of Fisheries, 7 p. (Available from Washington Department of Fish and Wildlife, 600 Capital Way N., Olympia WA 98501-1091.)
- Bugert, R.M., C. W. Hopley, C. A. Busack, and G. W. Mendel. 1995. Maintenance of stock integrity in Snake River fall chinook salmon. *Uses and Effects of Cultured Fishes in Aquatic Ecosystems*. American Fisheries Society, Bethesda, MD, Vol. 15: 267-276.
- Burgner, R. L. 1991. the life history of sockeye salmon (*Oncorhynchus nerka*). In C. Groot and L. Margolis (eds.), *Life history of Pacific salmon*, p. 3-117. Univ. B.C. Press, Vancouver, B.C.
- Busack, C. and D. Rawding. 2003. HPVA results for salmon and steelhead production in Washington Lower Columbia Basins. Appendix J in McElhany et al. Interim report on viability criteria for Willamette and Lower Columbia basin Pacific salmonids. NMFS-NWFSC.

- Carie, D. G. 2000. Spring and summer chinook salmon spawning ground surveys on the Entiat River, 2000. U.S. Fish & Wildlife Service Rept. Mid-Columbia River Fishery Resource Office. Leavenworth, Washington. 17 pp.
- CDFG (California Department of Fish and Game). 1998. Report to the Fish and Game Commission: a status review of the spring-run chinook salmon (*Oncorhynchus tshawytscha*) in the Sacramento River drainage. California Department of Fish and Game, Sacramento, CA.
- CDFG (California Department of Fish and Game). 2001. Final report on anadromous salmonid fish hatcheries in California. Technical report, California Department of Fish and Game and National Marine Fisheries Service Southwest Region. (<http://swr.nmfs.noaa.gov/HatcheryReviewPublicDraft2.pdf>)
- Campbell, B. 2000. Salmonid Stock Inventory (SaSSI) database.
- Castle, P. and N. Currens. 2001a. Personal communication from Nooksack comanagers to the Puget Sound Technical Recovery Team. December 11, 2001.
- Castle, P. and N. Currens. 2001b. North Fork Nooksack native spring chinook escapement methodology. Memo to the Puget Sound Technical Recovery Team. December 2001
- Chapman, D. W. 1986. Salmon and steelhead abundance in the Columbia River in the nineteenth century. Trans. Am. Fish. Soc. 115:662-670.
- Chilcote, M. W. 2001. Conservation Assessment of Steelhead Populations in Oregon. Oregon Department of Fish and Wildlife. Portland, OR.
- Cooney, T. 2001. Upper Columbia River steelhead and spring chinook salmon quantitative analysis report. Part 1: run reconstructions and preliminary analysis of extinction risks. National Marine Fisheries Service. Hydro Program. Technical Review Draft.
- Craig, J. A., and A. J. Suomela. 1941. Time of appearance of the runs of salmon and steelhead trout native to the Wenatchee, Entiat, Methow, and Okanogan Rivers. Cited In J. W. Mullan, K. R. Williams, G. Rhodus, T. W. Hillman, and J. D. McIntyre (editors). 1992. Production and habitat of salmonids in mid-Columbia River tributary streams. U.S. Fish Wildl. Serv. Monograph I:J358-J380.
- Cramer, D. 2002d. Sandy R Spring Chinook Counts at Marmot Dam. Portland General Electric Excel Workbook delivered via e-mail to Paul McElhany and Sarah Sydor, NWFSC.
- Cramer, D. 2002e. Portland General Electric - Clackamas River spring chinook adults. Portland General Electric. Data delivered via e-mail from Kathryn Kostow, Oregon Dept of Fish and Wildlife.

- Dauble, D. D., R. L. Johnson, and A. P. Garcia. 1999. [Fall chinook salmon spawning in the tailraces of lower Snake River hydroelectric projects](#). Trans. Amer. Fish Soc. 128 (4): 672-679.
- Dennis, B., P. L. Munholland, and J. M. Scott. 1991. Estimation of growth and extinction parameters for endangered species. Ecol. Mono. 61:115-143.
- FCRPS (Federal Columbia River Power System). 2000. Endangered Species Act - Section 7 Consultation, Biological Opinion; Reinitiation of Consultation on Operation of the Federal Columbia River Power System, Including the Juvenile Fish Transportation Program, and 19 Bureau of Reclamation Projects in the Columbia Basin between the U.S. Army Corps of Engineers, Bonneville Power Administration, Bureau of Reclamation, and the National Marine Fisheries Service. National Marine Fisheries Service, Northwest Region. December 21, 2000.
- Fisher, F. W. 1994. Past and present status of Central Valley chinook salmon. Conserv. Biol. 8:870-873.
- Ford, M, P. Budy, C. Busack, D. Chapman, T. Cooney, T. Fisher, J. Geiselman, T. Hillman, J. Lukas, C. Peven, C. Toole, E. Weber, and P. Wilson. 2001. Final report of the Upper Columbia River Steelhead and Spring Chinook Salmon Biological Requirements Committee, March 2001. National Marine Fisheries Service, Northwest Fisheries Science Center, Seattle, WA.
- Foster, C. A. 1998. Excerpt from: 1997 Willamette River spring chinook salmon run fisheries and passage at Willamette Falls (draft). Unpublished.
- Foster, C. A. 2000. 1999 Willamette River Spring Chinook Run, Fisheries, and Passage at Willamette Falls. (Draft).
- Foster, C. A. 2002. Willamette Falls Counts by Year (dynamic database).
- Fulop, J. 2002. Oregon Lower Columbia River Fall and Winter Chinook Spawning Ground Surveys, 1948-2000. Oregon Department of Fish and Wildlife, Columbia River Management. Portland, OR.
- Fulop, Jeff. 2003. Oregon Lower Columbia River Fall and Winter Chinook Spawning Ground Surveys, 1952-2002. Focus on 2001-2002 Oregon Department of Fish and Wildlife, Columbia River Management. Portland, OR.
- Fulton, L. A. 1968. Spawning areas and abundance of chinook salmon, *Oncorhynchus tshawytscha*, in the Columbia River Basin--Past and present. U.S. Fish. Wildl. Serv. Spec. Sci. Rep.--Fish. 571, 26 p.
- Gilbert, C. H. 1912. Age at maturity of Pacific coast salmon of the genus *Oncorhynchus*. Bull. U.S. Fish Comm. 32:57-70.

- Hallock, R. J. and F. W. Fisher. 1985. Status of winter-run chinook salmon, *Oncorhynchus tshawytscha*, in the Sacramento River. California Department of Fish and Game, Anadromous Fisheries Branch. 29 p.
- Hankin, D. G., J. W. Nicholas and T. W. Downey. 1993. Evidence for inheritance of age of maturity in chinook salmon, *Onchorhynchus tshawytscha*. Can. J. fish. Aquat. Sci.
- Hayman, B. 2002. Personal communication to Norma Jean Sands, NWFSC. January 2002.
- Healey, M. C. 1983. Coastwide distribution and ocean migration patterns of stream- and ocean type chinook salmon, *Oncorhynchus tshawytscha*. Canadian Field-Naturalist 97:427-433.
- Healey, M. C. 1986. Optimum size and age at maturity in Pacific salmon and effects of size selective fisheries. Can. Spec. Publ. Fish. Aquat. Sci. 89:39-52.
- Healey, M. C. 1991. The life history of chinook salmon (*Oncorhynchus tshawytscha*). In C. Groot and L. Margolis (eds), Life history of Pacific salmon, p. 311-393. Univ. BC Press, Vancouver, BC.
- Hedgecock, D. 2002. Microsatellite DNA for the management and protection of California's Central Valley chinook salmon (*Oncorhynchus tshawytscha*). Final report for the amendment to agreement No. B-59638. UC Davis, Bodega Bay, CA.
- Higgins P., S. Dobush, and D. Fuller. 1992. Factors in Northern California threatening stocks with extinction. Humboldt Chapter of the American Fisheries Society. 26 pp.
- Holmes, E. E. 2001. Estimating risks in declining populations with poor data. Proc. Nat. Acad. Sci. USA 98:5072-5077.
- Howell, P. J. 1986. Willamette basin fish management plan: status and progress 1979-85. ODFW, Portland, OR.
- Hubble, J. and S. Crampton. 2000. Methow basin spring chinook spawner ground survey report for 1999. Fisheries Resource Management Program. Yakama Nation. Prepared for Douglas County PUD. 17 pp + tables.
- Irving J. S. and T. Bjornn. 1991. A forecast of abundance of Snake River fall chinook salmon. Prepared for U.S. Fish & Wildlife Service. Unpubl. MS. 55 p. Available from Idaho Cooperative Fishery Research Unit. Univ. of Idaho, Moscow, ID.
- Johnson, T. 2003a. Age data (2001 and 2002), unpublished. March 27, 2003.
- Johnson, T. 2003b. Personal communication to Norma Jean Sands and co-managers. March 27, 2003.

- Joint Chinook Technical Committee (CTC). 1999. Annual reports for 1995 and 1996.
- Joint Chinook Technical Committee (CTC). 2000. CTC model and ER analyses output. Dell Simmons, December 2000.
- Kostow, K. 2002. Leaburg and McKenzie abundance from ODFW. Data delivered via e-mail, September 2002. Kreitman, G. 1981. Addendum to 8/13/80 naturally spawning population estimates memo. Streamnet Reference.
- Lampkis, N. 2003. Run reconstruction tables. Unpublished data. March 31, 2003.
- Lindley, S. T. and M. H. Mohr. 2003. Predicting the impact of striped bass (*Morone saxatilis*) population manipulations on the persistence of winter-run chinook salmon (*Oncorhynchus tshawytscha*). Fish. Bull., U.S. 101:321-331.
- Lindley, S. T. (in press). Estimation of population growth and extinction parameters from noisy data. *Ecol. Appl.*
- Marmorek, D. R., C. N. Peters and I. Parnell (eds). 1998. Plan for analyzing and testing hypotheses (PATH): final report for fiscal year 1998. ESSA Technologies Ltd., Vancouver, BC.
- Marshall, A. R., C. Smith, R. Brix, W. Dammers, J. Hymer, and L. LaVoy. 1995. Genetic diversity units and major ancestral lineages for chinook salmon in Washington. In C. Busack and J. B. Shaklee (eds.), Genetic diversity units and major ancestral lineages of salmonid fishes in Washington, p. 111-173. Wash. Dep. Fish Wildl. Tech. Rep. RAD 95-02. (Available from Washington Department of Fish and Wildlife, 600 Capital Way N., Olympia WA 98501-1091)
- Marshall, A. R., H. L. Blankenship, and W. P. Connor. 2000. Genetic Characterization of Naturally Spawned Snake River Fall-Run Chinook Salmon. Trans. Amer. Fish. Soc. 129: 680-698.
- Matthews, G. M., and R. S. Waples. 1991. Status review for Snake River spring and summer chinook salmon. NOAA Tech. Memo. NMFS F/NWC-200, 75 p. (Available from Natl. Mar. Fish. Serv., Northwest Fisheries Science Center, Coastal Zone and Estuarine Studies Division, 2725 Montlake Blvd. E., Seattle, WA 98112-2097.)
- McClure, M. M., E. E. Holmes, B. L. Sanderson, and C. E. Jordan. (in press). A large-scale, multi-species status assessment: anadromous salmonids in the Columbia River Basin. *Ecol. Appl.*
- McElhany, P., M. H. Ruckelshaus, M. J. Ford, T. C. Wainwright, and E. P. Bjorkstedt. 2000. Viable salmonid populations and the recovery of evolutionarily significant units. U. S. Dept. Commer. NMFS-NWFSC-42.

- McElhany, P., T. Backman, C. Busack, S. Heppell, S. Kolmes, A. Maule, J. Myers, D. Rawding, D. Shively, and C. Steward. 2002. Willamette/Lower Columbia Pacific salmonid viability criteria. Draft report from the Willamette/Lower Columbia Technical Recovery Team. December 2002.
- McElhany, P., T. Backman, C. Busack, S. Heppell, S. Kolmes, A. Maule, J. Myers, D. Rawding, D. Shively, A. Steel, and C. Steward. 2003. Interim report on viability criteria for Willamette and Lower Columbia basin Pacific salmonids. Report from the Willamette/Lower Columbia Technical Recovery Team. NMFS-NWFSC. March 2003.
- McPhail, J.D., and C. C. Lindsey. 1970. Freshwater fishes of Northwestern Canada and Alaska. *Bull. Fish. Res. Board Can.* 173:381.
- Miller, R. J., and E. L. Brannon. 1982. The origin and development of life-history patterns in Pacific salmon. *In* E. L. Brannon and E. O. Salo (eds), *Proceedings of the Salmon and Trout Migratory Behavior Symposium.*, p. 296-309. Univ. Wash. Press, Seattle, WA.
- Moffett, J. W. 1949. The first four years of king salmon maintenance below Shasta Dam, Sacramento River, California. *Cal. Fish Game* 35:77-102.
- Mosey, T. R. and L. J. Murphy. 2002. Spring and summer spawning ground surveys on the Wenatchee River basin, 2001. Washington Dept. of Fish & Wildlife Rept. to Chelan County PUD. 35 pp + appendices.
- Murtagh, T., J. Massey, and D. E. Bennett. 1997. Excerpt from: Sandy River basin fish management plan 1997 (draft). Unpublished. Streamnet reference.
- Myers, J. M., R. G. Kope, B. J. Bryant, D. Teel, L. J. Lierheimer, T. C. Wainwright, W. S. Grant, F. W. Waknitz, K. Neely, S. T. Lindley, and R. S. Waples. 1998. Status review of chinook salmon from Washington, Idaho, Oregon, and California. U.S. Dep. Commer., NOAA Tech. Memo NMFS-NWFSC-35, 443 p.
- Myers, J. M., C. Busack, D. Rawding, and A. Marshall. 2002. Identifying historical populations of chinook and chum salmon and steelhead within the lower Columbia River and upper Willamette River evolutionary significant units. Draft report to the co-managers from the Willamette/Lower Columbia River Technical Recovery Team (10 May 2002).
- Nehlsen W., J. E. Williams, and J. A. Lichatowich. 1991. Pacific salmon at the crossroads: stocks at risk from California, Oregon, Idaho, and Washington. *Fisheries* 16(2): 4-21.
- NMFS (National Marine Fisheries Service). 1987. Endangered and threatened species; winter-run chinook salmon. Federal Register [Docket No. 27 February 1986] 52(39):6041-6048.

- NMFS (National Marine Fisheries Service). 1995. Proposed recovery plan for Snake River Salmon. 364 p. + app. (Available from Environmental and Technical Services Division, Natl. Mar. Fish. Serv., 525 N.E. Oregon St., Suite 500, Portland, OR 97232.)
- NMFS (National Marine Fisheries Service). 1997. NMFS proposed recovery plan for the Sacramento River winter-run chinook. NMFS Southwest Region, Long Beach, CA.
- NMFS (National Marine Fisheries Service). 1998. Conclusions regarding the updated status of Puget Sound, Lower Columbia River, Upper Willamette River, and Upper Columbia River spring-run ESUs of West Coast Chinook Salmon. Memorandum to U. Varanasi (Northwest Fisheries Science Center, NMFS), W. Stelle (NWFSC, NMFS), and W. Hogarth (Southwest Fisheries Science Center, NMFS) from M. Schiewe (Northwest Fisheries Science Center, NMFS), 12 February 1999. 62 p.
- NMFS (National Marine Fisheries Service). 1999. Status review update for deferred ESUs of West Coast chinook salmon (*Oncorhynchus tshawytscha*) from Washington, Oregon, California, and Idaho. Memorandum to U. Varanasi (Northwest Fisheries Science Center, NMFS) and M. Tillman (Southwest Fisheries Science Center, NMFS) from M. Schiewe (Northwest Fisheries Science Center, NMFS), 16 July 1999. 116 p.
- NMFS (National Marine Fisheries Service). 2002. Meeting between Nooksack co-managers (Pete Castle and Dale Griggs) and NMFS (Susan Bishop and Norma Jean Sands). July 29, 2002.
- NMFS (National Marine Fisheries Service) and Co-managers. 2002. NMFS/Co-managers Meeting Point No Point. August 8, 2002.
- Nisqually Co-managers. 2002. Personal communication to the Puget Sound Technical Recovery Team.
- Norman, G. 1982. Population estimates of natural spawning adults and jack fall chinook on the Wind, Big White Salmon, and Klickitat rivers, 1964-81. Unpublished Streamnet Reference.
- NWIFC (Northwest Indian Fisheries Commission). 2001. Estimation of contribution of hatchery origin fall chinook salmon to Duwamish-Green River spawning ground populations. NWIFC memo to Distribution June 30, 2001.
- ODFW (Oregon Dept of Fish and Wildlife). 1990. Sandy River subbasin: Salmon and steelhead production plan. Oregon Department of Fish and Wildlife. Northwest Power Planning Council. Streamnet Reference.
- ODFW (Oregon Dept of Fish and Wildlife). 1998. ODFW Columbia River Management unpublished files: Estimated number of spawning fall chinook in lower Columbia River tributaries, 1964-1997. Unpublished abundance data and reference from Streamnet.org.

- ODFW (Oregon Department of Fish and Wildlife). 2002. Spring Chinook Salmon in the Willamette and Sandy Rivers. Oregon Department of Fish and Wildlife, Fish Research Project Oregon, Annual Progress Report. Portland, OR.
- ODFW (Oregon Dept of Fish and Wildlife). 1999. Field data forms and files of ODFW spawning ground surveys, Columbia River Management. ODFW.
- PFMC (Pacific Fishery Management Council). 1997. Review of the 1996 ocean salmon fisheries. PFMC, 275 p.
- PFMC (Pacific Fishery Management Council). 2001. Review of 2000 ocean salmon fisheries. (Document prepared for the Council and its advisory entities.). Technical report, Pacific Fishery Management Council, 7700 NE Ambassador Place, Suite 200, Portland, OR
- PFMC (Pacific Fishery Management Council). 2002a. Review of 2001 ocean salmon fisheries. (Document prepared for the Council and its advisory entities.). Technical report, Pacific Fishery Management Council, 7700 NE Ambassador Place, Suite 200, Portland, OR.
- PFMC (Pacific Fishery Management Council). 2002b. Preseason Report I Stock Abundance Analysis for 2002 Ocean Fisheries. Pacific Fishery Management Council, Portland, OR.
- PFMC (Pacific Fishery Management Council). Central Valley winter and spring chinook salmon workgroup report. Exhibit B.8.b, March 2003. Pacific Fishery Management Council, Portland, OR.
- PSC (Pacific Salmon Commission). 1994. Pacific Salmon Commission Joint Chinook Technical Committee 1993 annual report. Pacific Salmon Commission. Report Chinook (94)-1, 121 p. + app.
- PSC (Pacific Salmon Commission). 2002. Lower Columbia Harvest Rate. Unpublished. Excel spreadsheets provided by Dell Simmons that include age structure information. Based on PFMC Chinook Technical Committee CWT analysis.
- Phinney, C. 2001. Personal communication from Puyallup Indian Fisheries to Puget Sound Technical Recovery Team. January 25, 2001.
- Puget Sound Indian Tribes and WDFW (Washington Department of Fish and Wildlife). 2001. Puget Sound Comprehensive Chinook Management Plan: Harvest Management Component. March 2001.
- PSTRT (Puget Sound Technical Recovery Team). 2001. Independent populations of chinook salmon in Puget Sound. Puget Sound Technical Recovery Team Report. 92 pp.
- PSTRT (Puget Sound Technical Recovery Team). 2002. Planning ranges and preliminary guidelines for the delisting and recovery of the Puget Sound chinook salmon

- evolutionarily significant unit . Puget Sound Technical Recovery Team Report. 17 pp. (available at: http://www.nwfsc.noaa.gov/cbd/trt/trt_puget.htm).
- Rawding, D. 2001a. Lower Columbia Chinook Escapement (WDFW). Excel file provided by Dan Rawding in Fall 2001.
- Rawson, K. 2001. Terminal harvest rates for Snohomish River using terminal run reconstruction. November 2001.
- Rawson, K. and Kraemer, C. Personal communication to the Puget Sound Technical Recovery Team.
- Rawson, K. and Kraemer, C. 2001. Unpublished data from otolith sampling on spawning grounds. February 2001.
- Reynolds, F. L., T. J. Mills, R. Benthin, and A. Low. 1993. Restoring Central Valley streams: a plan for action. California Department of Fish and Game, Sacramento, CA.
- SSHAG (Salmon Steelhead Hatchery Assessment Group). 2003. Hatchery Broodstock Summaries and Assessments for Chum Salmon, Coho Salmon, Chinook Salmon, and Steelhead within Listed and ESUs. Technical review draft. NMFS Northwest and Southwest Fisheries Science Centers.
- Sands, N. J. A User's Guide to the abundance and productivity tables (Excel Workbook). In prep.
- Sanford, B. 2003. WDFW pers. com. Washington chinook escapement data emailed to Paul McElhany April 1, 2003.
- Scott, J. 2002. Personal communication to the Puget Sound Technical Recovery Team. May 21, 2002.
- Serl, J. and Morrill C. 2002. Draft 2002 annual report for the Cowlitz Falls project. Prepared by WDFW for BPA contract number 96B192557.
- Shardlow, T.F., T. M. Webb and D. T. Lightly. 1986. Chinook salmon escapement estimation on the Campbell and Quinsam Rivers in 1984: accuracy and precision of mark/recapture techniques using tagged salmon carcasses. Canadian Technical Report of Fisheries and Aquatic Sciences 1507. 52 p.
- Snider, B. and B. Reavis and S. Hill. 1999. 1998 Upper Sacramento River winter-run chinook salmon escapement survey May-August 1998. California Department of Fish and Game, Water and Aquatic Habitat Conservation Branch, Stream Evaluation Program Report. January 1999.
- Steel, E. A. and M. B. Sheer. 2002. Broad-scale habitat analyses to estimate fish densities for viability criteria. Appendix C in Willamette/Lower Columbia Pacific salmonid viability criteria. Willamette/Lower Columbia Technical Recovery Team Report. December 2002.

- Steel, A. and M. Sheer. 2003. Broad-scale habitat analysis to estimate fish densities for viability criteria. Appendix I in McElhany et al. Interim report on viability criteria for Willamette and lower Columbia basin Pacific salmonids. NMFS-NWFSC.
- USFWS (United States Fish and Wildlife Service). 1997. Documents submitted to the ESA Administrative Record for west coast chinook salmon by D. Finberg, 26 February 1997, 4p.
- Wahle R. J., Pearson. 1987. A listing of Pacific Coast spawning streams and hatcheries producing chinook and coho spawners. NOAA Tech. Memo. NMFS F/NWC-122, 32 p .
- Waknitz, F. W., G. M. Matthews, T. Wainwright, and G. A. Winans. 1995. Status review for Mid-Columbia River summer chinook salmon. NOAA Tech. Mem. NMFS-NWFSC-22, 80 p.
- Waples, R. S. 1991. Pacific salmon, *Oncorhynchus* spp., and the definition of "species" under the Endangered Species Act. Marine Fisheries Review 53:11-22.
- Waples, R. S., J. Robert, P. Jones, B. R. Beckman, and G. A. Swan. 1991. Status review for Snake River fall chinook salmon. NOAA Tech. Memo. NMFS F/NWC-201, 73 p. (Natl. Mar. Fish. Serv., Northwest Fisheries Science Center, Coastal Zone and Estuarine Studies Division, 2725 Montlake Blvd. E., Seattle, WA 98112-2097.)
- Ward, P., T. McReynolds, and C. Garman. 2002. Butte and Big Chico Creeks spring-run chinook salmon, *Oncorhynchus Tshawytscha*, life history investigation 2000-2001. California Department of Fish and Game, Chico, CA.
- WDF (Washington Department of Fisheries), Washington Department of Wildlife (WDW), and Western Washington Treaty Indian Tribes (WWTIT). 1993. 1992 Washington State salmon and steelhead stock inventory (SASSI). Wash. Dep. Fish Wildl., Olympia, 212 p. + 5 regional volumes. (Available from Washington Department of Fish and Wildlife, 600 Capitol Way N., Olympia, WA 98501-1091.)
- WDFW (Washington Department of Fish and Wildlife) and Point No Point Treaty Tribes. 2000. Summer Chum Salmon Conservation Initiative: An implementation plan to recover summer chum salmon in the Hood Canal and Strait of Juan de Fuca Region. April 2000.
- WDFW (Washington Department of Fish and Wildlife) et al. 2001a. Data obtained from the WDFW age database. March 2001.
- WDFW (Washington Department of Fish and Wildlife) and Point No Point Treaty Tribes. 2001b. Summer Chum Salmon Conservation Initiative. Supplemental Report No. 3 Annual Report for the 2000 summer chum salmon return to the Hood Canal and Strait of Juan de Fuca Region. December 2001.

- WDFW (Washington Department of Fish and Wildlife), Point No Point and Makah Tribes. 2001c. Management Framework Plan and Salmon Runs' Status for the Strait of Juan de Fuca. November 2001.
- Yoshiyama, R. M., E. R. Gerstung, F. W. Fisher, and P. B. Moyle. 2001. Historic and present distribution of chinook salmon in the Central Valley drainage of California. In R. L. Brown, editor, *Fish Bulletin 179: Contributions to the biology of Central Valley salmonids.*, volume 1, pages 71-176. California Department of Fish and Game, Sacramento, CA.